

Global Existence for the Vlasov-Poisson System with Steady Spatial Asymptotics

Stephen Pankavich
 Department of Mathematical Sciences
 Carnegie Mellon University
 Pittsburgh, PA 15213
 sdp@andrew.cmu.edu

January 5, 2010

MOS Classification : 35L60, 35Q99, 82C21, 82C22, 82D10.

Keywords : Vlasov, kinetic theory, Cauchy problem, global existence, spatial decay.

Abstract

A collisionless plasma is modelled by the Vlasov-Poisson system in three space dimensions. A fixed background of positive charge, dependant upon only velocity, is assumed. The situation in which mobile negative ions balance the positive charge as $|x| \rightarrow \infty$ is considered. Thus, the total positive charge and the total negative charge are both infinite. Smooth solutions with appropriate asymptotic behavior for large $|x|$, which were previously shown to exist locally in time, are continued globally. This is done by showing that the charge density decays at least as fast as $|x|^{-6}$. This paper also establishes decay estimates for the electrostatic field and its derivatives.

INTRODUCTION

Let $F : \mathbb{R}^3 \rightarrow [0, \infty)$, $f_0 : \mathbb{R}^3 \times \mathbb{R}^3 \rightarrow [0, \infty)$, and $A : [0, \infty) \times \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be given. We seek a solution, $f : [0, \infty) \times \mathbb{R}^3 \times \mathbb{R}^3 \rightarrow [0, \infty)$ satisfying

$$\left. \begin{aligned} \partial_t f + v \cdot \nabla_x f - (E + A) \cdot \nabla_v f &= 0, \\ \rho(t, x) &= \int (F(v) - f(t, x, v)) dv, \\ E(t, x) &= \int \rho(t, y) \frac{x-y}{|x-y|^3} dy, \\ f(0, x, v) &= f_0(x, v). \end{aligned} \right\} \quad (1)$$

Here F describes a number density of positive ions which form a fixed background, and f denotes the density of mobile negative ions in phase space. Notice that if $f_0(x, v) = F(v)$ and $A = 0$, then $f(t, x, v) = F(v)$ is a steady solution. Thus, we seek solutions for which $f(t, x, v) \rightarrow F(v)$ as $|x| \rightarrow \infty$. It is important to notice that (1) is a representative problem, and that problems concerning multiple species of ions can be treated in a similar manner.

Precise conditions which ensure local existence and conditions for continuation were given in [18]. Also, *a priori* bounds on ρ and a quantity related to the energy were obtained in [17]. Therefore, we will work towards establishing global existence with these bounds and using similar assumptions. Other work on the infinite mass case has been done in [9] and [3].

The case when $F(v) = 0$ and $f \rightarrow 0$ as $|x| \rightarrow \infty$ has been studied extensively. Smooth solutions were shown to exist globally in time in [14] and independently in [11]. Important results prior to global existence appear in [1], [5], [6], [7], and [10]. Also, the method used by [14] is refined in [8] and [16]. Global existence for the Vlasov-Poisson system in two dimensions was established in [12] and [19]. A complete discussion of the literature concerning Vlasov-Poisson may be found in [4]. We also mention [2] and [15] since the problem treated in these papers is periodic in space, and thus the solution does not decay for large $|x|$.

1 Section 1

Let $p \in (3, 4)$ and denote

$$R(x) = R(|x|) = (1 + |x|^2)^{\frac{1}{2}}.$$

We will use the following notation :

$$\|g\|_{L^\infty(\mathbb{R}^n)} = \sup_{z \in \mathbb{R}^n} |g(z)|$$

and

$$\|\rho\|_p := \|\rho(x) R^p(x)\|_{L^\infty(\mathbb{R}^3)},$$

but never use L^p . We will write, for example, $\|\rho(t)\|_p$ for the $\|\cdot\|_p$ norm of $x \mapsto \rho(t, x)$.

Following [17] and [18], we assume the following conditions hold for some $C > 0$ and all $t \geq 0$, $x \in \mathbb{R}^3$, and $v \in \mathbb{R}^3$, unless otherwise stated :

(I) $F(v) = F_R(|v|)$ is nonnegative and C^2 with

$$F_R''(0) < 0, \quad (2)$$

and there is $W \in (0, \infty)$ such that

$$\left. \begin{aligned} F_R'(u) &< 0 & \text{for } u \in (0, W) \\ F_R(u) &= 0 & \text{for } u \geq W. \end{aligned} \right\} \quad (3)$$

(II) f_0 is C^1 and nonnegative.

(III) A is C^1 with

$$|A(t, x)| \leq C^{(0)} R^{-2}(x), \quad (4)$$

$$|\partial_{x_i} A(t, x)| \leq C R^{-3}(x), \quad (5)$$

and

$$\nabla_x \cdot A(t, x) = 0. \quad (6)$$

Finally, we assume there is a continuous function $a : [0, T] \rightarrow \mathbb{R}$ such that

$$\left| A(t, x) - a(t) \frac{x}{|x|^3} \right| \leq C R^{1-p}(x).$$

(IV) $F - f_0$ has compact support in v , and there is $N > 0$ such that for $|x| > N$, we have

$$|F(v) - f_0(x, v)| \leq C R^{-6}(x).$$

Then, we have global existence:

Theorem 1 *Assuming conditions (I)-(IV) hold, there exists $f \in \mathcal{C}^1([0, \infty) \times \mathbb{R}^3 \times \mathbb{R}^3)$ that satisfies (1) with $\|\int (F - f)(t) dv\|_p$ bounded on $t \in [0, T]$, for every $T > 0$. Moreover, f is unique.*

In addition, due to the previously known result of [13], stated here as Theorem 2, we are able to conclude further decay of the charge density in Corollary 1 :

Theorem 2 *Let $T > 0$ and f be the $\mathcal{C}^1$ solution of (1) on $[0, T] \times \mathbb{R}^3 \times \mathbb{R}^3$. Then, we have*

$$\|\rho(t)\|_6 \leq C_{p,t}$$

for any $t \in [0, T]$, where $C_{p,t}$ depends upon

$$\sup_{\tau \in [0, t]} \|\rho(\tau)\|_p.$$

Corollary 1 *Let $T > 0$ and f be the $\mathcal{C}^1$ solution of (1) on $[0, T] \times \mathbb{R}^3 \times \mathbb{R}^3$. Then, we have*

$$\left\| \int (F - f)(t) \, dv \right\|_6 \leq C$$

for any $t \in [0, T]$.

To prove Theorem 1, we will use Theorems 1 and 3 of [18], which guarantee local existence and continuation of the local solution so long as $\|\int (F - f)(t)\|_p$ is bounded for some $p > 3$. Therefore, the following lemma will be all that is needed to complete the proof of the theorem.

Lemma 1 *Assume conditions (I)-(IV) hold. Let f be a $\mathcal{C}^1$ solution of (1) on $[0, T] \times \mathbb{R}^3 \times \mathbb{R}^3$. Then,*

$$\left\| \int (F - f)(t) \, dv \right\|_p \leq C$$

for all $t \in [0, T]$, where C is determined by F , A , f_0 , and T .

Define

$$Q_f(t) := \sup\{|v| : \exists x \in \mathbb{R}^3, \tau \in [0, t] \text{ such that } f(\tau, x, v) \neq 0\} \quad (7)$$

and

$$Q_g(t) := \max\{W, Q_f(t)\}.$$

In Section 2, we will bound E , ∇E , and $\nabla_{x,v} f$. Also, we will estimate the energy, and obtain bounds on $Q_f(t)$ and thus $Q_g(t)$. To better reveal the line of thought, the proofs of Lemmas 2 through 4 are deferred to Section 3.

We will denote by “ C ” a generic constant which changes from line to line and may depend upon f_0 , A , F , or T , but not on t , x , or v . When it is necessary to refer to a specific constant, we will use numeric superscripts to distinguish them. For example, $C^{(0)}$, as in (III), will always refer to the same numerical constant.

To estimate E and ∇E , we will use

Lemma 2 *For any $q > 0$ and $b \in [0, \frac{5}{18})$ with $b \leq \frac{2}{q}$, we have*

$$|E(t, x)| \leq C (\|\rho(t)\|_q R^{-q}(x))^b$$

for any $t \in [0, T]$, $|x| \geq 1$, where C may depend upon $\|\rho(t)\|_\infty$.

and

Lemma 3 *For any $q > 0$ and $a \in [0, 1)$ with $a \leq \frac{3}{q}$, we have*

$$|\nabla E(t, x)| \leq C (\|\rho(t)\|_q R^{-q}(x))^a$$

for any $t \in [0, T]$, $|x| \geq 1$, where C may depend upon $\|\rho(t)\|_\infty$ and $\|\nabla \rho(t)\|_\infty$.

Then, we will use the following lemma to bound the p -norm of ρ and obtain decay of both E and ∇E :

Lemma 4 *For any $q \in [0, \frac{54}{13})$, $\|\rho(t)\|_q$ is bounded for $t \in [0, T]$, where C may depend upon $\|\rho(t)\|_\infty$, $\|\nabla \rho(t)\|_\infty$, and $Q_f(T)$.*

Thus, once we show $\|\rho(t)\|_\infty$ and $\|\nabla \rho(t)\|_\infty$ are bounded for $t \in [0, T]$, and $Q_f(T)$ is finite, we may use Lemma 4 to bound $\|\rho(t)\|_p$ since $p \in (3, 4)$. Then, we can prove Lemma 1, and thus Theorem 1.

2 Section 2

2.1 Characteristics

Define the characteristics, $X(s, t, x, v)$ and $V(s, t, x, v)$, by

$$\left. \begin{aligned} \frac{\partial X}{\partial s}(s, t, x, v) &= V(s, t, x, v) \\ \frac{\partial V}{\partial s}(s, t, x, v) &= -(E(s, X(s, t, x, v)) + A(s, X(s, t, x, v))) \\ X(t, t, x, v) &= x \\ V(t, t, x, v) &= v. \end{aligned} \right\} \quad (8)$$

Then, we have

$$\frac{\partial}{\partial s} f(s, X(s, t, x, v), V(s, t, x, v)) = \partial_t f + V \cdot \nabla_x f - (E + A) \cdot \nabla_v f = 0.$$

Therefore, f is constant along characteristics, and

$$f(t, x, v) = f(0, X(0, t, x, v), V(0, t, x, v)) = f_0(X(0, t, x, v), V(0, t, x, v)). \quad (9)$$

Thus, we find by (II) that f is nonnegative.

Define $g : [0, \infty) \times \mathbb{R}^3 \times \mathbb{R}^3 \rightarrow \mathbb{R}$ by

$$g(t, x, v) := F(v) - f(t, x, v).$$

Then, we see that $\sup_{x,v} |g| \leq \|F\|_{L^\infty} + \|f_0\|_{L^\infty} < \infty$, and

$$\begin{aligned} \frac{\partial}{\partial s} g(s, X(s, t, x, v), V(s, t, x, v)) &= \partial_t g + V \cdot \nabla_x g - (E + A) \cdot \nabla_v g \\ &= -\nabla_v F(V(s)) \cdot (E + A)(s, X(s)). \end{aligned} \quad (10)$$

Finally, for any $s \in [0, T]$ with $f(s, X(s), V(s)) \neq 0$, we have for $t \in [0, T]$ and $x, v \in \mathbb{R}^3$,

$$\begin{aligned} |X(s, t, x, v) - x| &= \left| \int_s^t \dot{X}(\tau, t, x, v) \, d\tau \right| \\ &\leq \int_s^t |V(\tau, t, x, v)| \, d\tau \\ &\leq TQ_g(T). \end{aligned}$$

So, assuming we can bound $Q_g(T)$, we have for $|x| \geq 2TQ_g(T)$, $v \in \mathbb{R}^3$, $t \in [0, T]$, and $s \in [0, t]$,

$$|X(s, t, x, v)| \geq |x| - TQ_g(T) \geq \frac{1}{2}|x| \quad (11)$$

and

$$|X(s, t, x, v)| \leq |x| + TQ_g(T) \leq \frac{3}{2}|x|. \quad (12)$$

Unless it is necessary, we will omit writing the dependence of $X(s)$ and $V(s)$ on t , x , and v for the remainder of the paper.

2.2 Bounds on the Field

Much of the work that follows will rely on energy estimates found in [17]. In particular, we combine Lemma 3 and Theorem 4 from [17] to obtain an a priori bound on ρ .

First, define $\sigma : [0, \infty) \rightarrow \mathbb{R}$ by

$$\sigma(h) = - \int_0^{\min\{h, F(0)\}} \left(F^{-1}(\tilde{h}) \right)^2 d\tilde{h}$$

and $S : [0, \infty) \times [0, \infty) \rightarrow [0, \infty)$ by

$$S(h, \eta) = (h - F(\eta))\eta^2 + \sigma(h) - \sigma(F(\eta)).$$

Then, from [17], we have the following lemma:

Lemma 5 *Let*

$$k(t, x) = \int S(f(t, x, v), |v|) dv.$$

Assuming conditions (I) – (IV) hold, we have

$$\int |F(v) - f(t, x, v)| dv \leq C(k(t, x)^{\frac{3}{5}} + k(t, x)^{\frac{1}{2}}) \quad (13)$$

and

$$\int k(t, x) dx \leq C. \quad (14)$$

It follows directly from (13) that

$$|\rho(t, x)| \leq C(k(t, x))^{\frac{3}{5}} + k(t, x)^{\frac{1}{2}} \quad (15)$$

Then, using Lemma 5 , we may bound the field.

Assume $\|\rho(t)\|_{L^\infty(\mathbb{R}^6)} \leq C$ for all $t \in [0, T]$. Then, for $t \geq 0$, $x \in \mathbb{R}^3$, and any $R > 0$

$$\left. \begin{aligned} |E(t, x)| &\leq \int \frac{|\rho(t, y)|}{|x-y|^2} dy \\ &\leq \int_{|x-y| < R} \|\rho(t)\|_{L^\infty} |x-y|^{-2} dy + \int_{|x-y| > R} |\rho(t, y)| |x-y|^{-2} dy \\ &\leq 4\pi \|\rho(t)\|_{L^\infty} \int_0^R dr + C \int_{|x-y| > R} (k^{\frac{1}{2}}(t, y) + k^{\frac{3}{5}}(t, y)) |x-y|^{-2} dy \\ &\leq 4\pi \|\rho(t)\|_{L^\infty} R + C \left(\int k(t, y) dy \right)^{\frac{1}{2}} \left(\int_{|x-y| > R} |x-y|^{-4} dy \right)^{\frac{1}{2}} \\ &\quad + C \left(\int k(t, y) dy \right)^{\frac{3}{5}} \left(\int_{|x-y| > R} |x-y|^{-5} dy \right)^{\frac{2}{5}} \\ &\leq 4\pi \|\rho(t)\|_{L^\infty} R + C \left(\int_R^\infty r^{-2} dr \right)^{\frac{1}{2}} + C \left(\int_R^\infty r^{-3} dr \right)^{\frac{2}{5}} \\ &\leq C(\|\rho(t)\|_{L^\infty} R + R^{-\frac{1}{2}} + R^{-\frac{4}{5}}). \end{aligned} \right\} \quad (16)$$

Obviously, we may choose $R = 1$ and deduce that $|E(t, x)| \leq C$. Suppose that the v -support of g is bounded for bounded times. Then, we find

$$|\rho(t, x)| \leq \int_{|v| \leq Q_g(t)} |g(t, x, v)| dv \leq (\|f_0\|_{L^\infty} + \|F\|_{L^\infty}) Q_g^3(t) \leq C Q_g^3(t).$$

Thus, we know for every $t \in [0, T]$,

$$\|\rho(t)\|_{L^\infty} \leq C Q_g^3(t).$$

Now, choose $R = Q_g^{-\frac{5}{3}}(t)$. If $R \geq 1$, then $Q_g(t)$ is bounded. Otherwise, $R \leq 1$ and thus $R^{-\frac{1}{2}} \leq R^{-\frac{4}{5}}$.

Then, we have

$$\left. \begin{aligned} |E(t, x)| &\leq C(\|\rho(t)\|_{L^\infty} R + R^{-\frac{4}{5}}) \\ &\leq C \left(Q_g^3(t) Q_g^{-\frac{5}{3}}(t) + (Q_g^{-\frac{5}{3}}(t))^{-\frac{4}{5}} \right) \\ &\leq C Q_g^{\frac{4}{3}}(t) \\ &=: C^{(1)} Q_g^{\frac{4}{3}}(t) \end{aligned} \right\} \quad (17)$$

2.3 Bounds on Derivatives of Density and Field

We proceed as in Section 4.2.5 of [4] with one modification : we will not assume that $\rho(t) \in L^1(\mathbb{R}^3)$ for all $t \in [0, T]$. Instead, we find for any $R > 0$, (letting $r = |x - y|$)

$$\begin{aligned} \left| \frac{1}{4\pi} \int_{|y-x| \geq R} \rho(t, y) \left(\frac{3(y_k - x_k)^2}{r^5} - \frac{1}{r^3} \right) dy \right| &\leq \frac{1}{\pi} \int_{|y-x| \geq R} |\rho(t, y)| r^{-3} dy \\ &\leq C \int_{|y-x| \geq R} \left(k^{\frac{1}{2}}(t, y) + k^{\frac{3}{5}}(t, y) \right) r^{-3} dy \\ &\leq C \left(\left(\int k(t, y) dy \right)^{\frac{1}{2}} \left(\int_{|y-x| \geq R} r^{-6} dy \right)^{\frac{1}{2}} \right. \\ &\quad \left. + \left(\int k(t, y) dy \right)^{\frac{3}{5}} \left(\int_{|y-x| \geq R} r^{-\frac{15}{2}} dy \right)^{\frac{2}{5}} \right) \\ &\leq C \left(R^{-\frac{3}{2}} + R^{-\frac{9}{5}} \right). \end{aligned}$$

Thus, introducing this estimate into the result of [4], we have for any $0 < d \leq R$

$$|\nabla_x E(t, x)| \leq C(1 + \ln(R/d)) \|\rho(t)\|_{L^\infty} + Cd \|\nabla_x \rho(t)\|_{L^\infty} + C(R^{-\frac{3}{2}} + R^{-\frac{9}{5}}). \quad (18)$$

Then, we may follow the argument in Section 4.2.6 of [4] and find a priori bounds on $\|\nabla_x E(t)\|_{L^\infty}$ and $\|\nabla_{x,v} f(t)\|_{L^\infty}$ as long as $\|\rho(t)\|_{L^\infty}$ is bounded. This work will be included in Appendix A.

2.4 Energy Bound

First, notice from (3) that we may write $W > 0$ as

$$W = \inf\{\eta > 0 : F(\eta) = 0\}.$$

Recall the definitions of σ , S , and k , as well as, (14), which we may write as

$$\int \int S(f(t, x, v), |v|) dv dx \leq C$$

for any $t \in [0, T]$.

Let $\eta \geq 2W$. We find

$$\begin{aligned} S(h, \eta) &= (h - F(\eta))\eta^2 + \sigma(h) - \sigma(F(\eta)) \\ &= h\eta^2 + \sigma(h) - \sigma(0) \\ &= h\eta^2 + \sigma(h) \\ &= h\eta^2 - \int_0^{\min\{h, F(0)\}} (F^{-1}(\tilde{h}))^2 d\tilde{h} \\ &\geq h\eta^2 - \min\{h, F(0)\}(F^{-1}(0))^2 \\ &\geq h\eta^2 - hW^2 \\ &= h(\eta^2 - W^2). \end{aligned}$$

Then, since $\eta \geq 2W$, we have $\frac{1}{4}\eta^2 \geq W^2$ and

$$\begin{aligned} h(\eta^2 - W^2) &= \frac{1}{2}h\eta^2 + h(\frac{1}{2}\eta^2 - W^2) \\ &\geq \frac{1}{2}h\eta^2. \end{aligned}$$

Thus, for any $h \geq 0$ and $\eta \geq 2W$,

$$S(h, \eta) \geq \frac{1}{2}h\eta^2$$

and, finally, for $P \geq 2W$

$$\int \int_{|v| > P} |v|^2 f(t, x, v) dv dx \leq 2 \int \int S(f(t, x, v), |v|) dv dx \leq C. \quad (19)$$

2.5 The Good, the Bad, and the Ugly Revisited

The method we will employ is very similar to that used in Section 4.4 of [4]. The differences are due mainly to the lack of positivity in the charge density, changes in the conserved energy, and the contribution of the applied field. We will assume throughout this section that $Q_g(t)$ is not already bounded and (17) applies.

Define

$$Q(t) := \left(\max\{(2W)^{\frac{4}{3}}, C^{(0)}\} \right)^{\frac{15}{13}} + Q_f(t).$$

Then, define $C^{(2)} := (C^{(0)})^{-\frac{7}{13}} + C^{(1)}$ so that we use (17) and (III) to find

$$\begin{aligned}
|E(\tau, x) + A(\tau, x)| &\leq C^{(1)} Q^{\frac{4}{3}}(t) + C^{(0)} R^{-2}(x) \\
&\leq C^{(1)} Q^{\frac{4}{3}}(t) + (C^{(0)})^{-\frac{7}{13}} (C^{(0)})^{\frac{20}{13}} \\
&\leq C^{(1)} Q^{\frac{4}{3}}(t) + (C^{(0)})^{-\frac{7}{13}} Q^{\frac{4}{3}}(t) \\
&= C^{(2)} Q^{\frac{4}{3}}(t)
\end{aligned}$$

for any $\tau \in [0, t]$.

Now, let $(\hat{X}(t), \hat{V}(t))$ be any fixed characteristic :

$$\frac{d}{dt} \hat{X} = \hat{V}, \frac{d}{dt} \hat{V} = E(t, \hat{X})$$

for which

$$f(t, \hat{X}(t), \hat{V}(t)) \neq 0.$$

For any $0 \leq \Delta \leq t$, we have

$$\int_{t-\Delta}^t |E(s, \hat{X}(s))| ds \leq C \int_{t-\Delta}^t \int \int \frac{|g(s, y, w)|}{|y - \hat{X}(s)|^2} dw dy ds \quad (20)$$

$$= C \int_{t-\Delta}^t \int \int \frac{|g(s, X(s, t, x, v), V(s, t, x, v))|}{|X(s, t, x, v) - \hat{X}(s)|^2} dv dx ds. \quad (21)$$

Let $P = Q^{\frac{13}{20}}(t)$, $R > 0$, and $\Delta = \frac{P}{4C^{(2)}Q^{\frac{4}{3}}(t)}$. From the definition of Q , notice that $P \geq 2W$.

Let us partition the integral into I_G , I_B , and I_U , where I_A is the integral in (21) over the set A , and the three sets are defined as :

$$G := \{(s, x, v) : t - \Delta < s < t \text{ and } (|v| < P \text{ or } |v - \hat{V}(t)| < P)\},$$

$$B := \{(s, x, v) : t - \Delta < s < t \text{ and } |v| > P \text{ and } |v - \hat{V}(t)| > P \\ \text{and } |X(s, t, x, v) - \hat{X}(s)| < R\},$$

$$U := \{(s, x, v) : t - \Delta < s < t \text{ and } |v| > P \text{ and } |v - \hat{V}(t)| > P \\ \text{and } |X(s, t, x, v) - \hat{X}(s)| > R\}.$$

We will use the invertibility of the characteristics as described in [4], so that when we set

$$y = X(s, t, x, v)$$

$$w = V(s, t, x, v)$$

we can invert using

$$x = X(t, s, y, w)$$

$$v = V(t, s, y, w).$$

In particular, notice $w = V(s, t, X(t, s, y, w), V(t, s, y, w))$ and $\frac{\partial(y, w)}{\partial(x, v)} = 1$.

To handle the integral over G , we must first deal with some preliminary inequalities :

1. First notice that $|V(s, t, x, v) - v| \leq \int_s^t |E(\tau, X(\tau)) + A(\tau, X(\tau))| d\tau$.

Thus, for $s \in [t - \Delta, t]$, we have

$$|V(s, t, x, v) - v| \leq \Delta C^{(2)} Q^{\frac{4}{3}}(t) = \frac{1}{4}P.$$

2. For $|v| < P$,

$$|V(s, t, x, v)| \leq |v| + \frac{1}{4}P < 2P.$$

3. For $|v - \hat{V}(t)| < P$,

$$|V(s, t, x, v) - \hat{V}(s)| \leq |v - \hat{V}(t)| + |\hat{V}(t) - \hat{V}(s)| + |V(s, t, x, v) - v| \leq P + \frac{1}{4}P + \frac{1}{4}P < 2P.$$

4. For $|v| > P$ and $\tau \in [t - \Delta, t]$,

$$\begin{aligned} |V(\tau, t, x, v)| &\geq |v| - \frac{1}{4}P \\ &> \frac{3}{4}P \\ &\geq \frac{3}{4} \cdot 2W \\ &> W \end{aligned}$$

Now, let

$$\chi_G(s, x, v) := \begin{cases} 1 & (s, x, v) \in G \\ 0 & \text{else.} \end{cases}$$

Then, we have

$$\begin{aligned}
I_G &= \int \int_G \int \frac{|g(s, X(s, t, x, v), V(s, t, x, v))|}{|X(s, t, x, v) - \hat{X}(s)|^2} dv dx ds \\
&= \int_{t-\Delta}^t \int \int \frac{\chi_G(s, x, v) |g(s, X(s, t, x, v), V(s, t, x, v))|}{|X(s, t, x, v) - \hat{X}(s)|^2} dv dx ds \\
&= \int_{t-\Delta}^t \int \int \frac{\chi_G(s, X(t, s, y, w), V(t, s, y, w)) |g(s, y, w)|}{|y - \hat{X}(s)|^2} dw dy ds.
\end{aligned}$$

If $\chi_G \neq 0$, then $|V(t, s, y, w)| < P$ or $|V(t, s, y, w) - \hat{V}(t)| < P$. Then, by Preliminary Inequalities 2 and 3, we have either $|w| < 2P$ or $|w - \hat{V}(s)| < 2P$. Set

$$\tilde{\rho}(s, y) := \int |g(s, y, w)| \chi_G(s, X(t, s, y, w), V(t, s, y, w)) dw.$$

Then, $\|\tilde{\rho}(s)\|_{L^\infty} \leq CP^3$. Also, using (13), we know

$$\tilde{\rho}(s, y) \leq \int |g(s, y, w)| dw \leq C(k^{\frac{3}{5}}(s, y) + k^{\frac{1}{2}}(s, y)).$$

Thus, we employ the method of (16) and (17) to find

$$\int \frac{\tilde{\rho}(s, y)}{|y - \hat{X}(s)|^2} dy \leq CP^{\frac{4}{3}}$$

and, finally, we have

$$I_G = \int_{t-\Delta}^t \int \frac{\tilde{\rho}(s, y)}{|y - \hat{X}(s)|^2} dy ds \leq C\Delta P^{\frac{4}{3}}. \quad (22)$$

Estimating I_B , we have

$$\begin{aligned}
I_B &= \int \int_B \int \frac{|g(s, X(s, t, x, v), V(s, t, x, v))|}{|X(s, t, x, v) - \hat{X}(s)|^2} dv dx ds \\
&\leq \int_{t-\Delta}^t \int_{|y - \hat{X}(s)| < R} \int \frac{|g(s, y, w)|}{|y - \hat{X}(s)|^2} dw dy ds \\
&\leq (\|F\|_{L^\infty} + \|f_0\|_{L^\infty}) Q^3(t) \int_{t-\Delta}^t \int_{|y - \hat{X}(s)| < R} |y - \hat{X}(s)|^{-2} dw ds \\
&\leq CQ^3(t) \int_{t-\Delta}^t \int_0^R dr ds,
\end{aligned}$$

and thus

$$I_B \leq C\Delta Q^3(t)R. \quad (23)$$

Finally, to estimate I_U , we use Section 3 (specifically, line (15)) of [16] to find

$$\int_{t-\Delta}^t |X(s, t, x, v) - \hat{X}(s)|^{-2} \chi_U(s, x, v) \, ds \leq \frac{C}{RP}, \quad (24)$$

for (x, v) as in U . The proof of this result is quite long, so we shall include a sketch rather than the entire proof. First, let

$$Z(s) = X(s, t, x, v) - \hat{X}(s).$$

Then, choose $s_0 \in [t - \Delta, t]$ such that

$$|Z(s_0)| \leq |Z(s)|$$

for all $s \in [t - \Delta, t]$. It is shown in [16] that

$$|Z(s)| \geq \frac{1}{4}P|s - s_0|. \quad (25)$$

Now, define

$$\Sigma(r) := \begin{cases} \frac{1}{R^2} & 0 \leq r \leq R^2 \\ \frac{1}{r} & r \geq R^2. \end{cases}$$

Notice that Σ is non-negative, non-increasing and

$$|Z(s)|^{-2} \chi_U(s, x, v) \leq \Sigma(|Z(s)|^2).$$

Using these properties of Σ with (25), we find

$$\begin{aligned} \int_{t-\Delta}^t |Z(s)|^{-2} \chi_U(s, x, v) \, ds &\leq \int_{t-\Delta}^t \Sigma(|Z(s)|^2) \, ds \\ &\leq \int_{t-\Delta}^t \Sigma\left(\left(\frac{1}{4}P|s - s_0|\right)^2\right) \, ds \\ &\leq \int \Sigma\left(\frac{P^2}{16}\tau^2\right) \, d\tau \\ &= \frac{16}{PR}. \end{aligned}$$

This shows (24).

Now, using (24) with (3), (9), (19), and preliminary inequality 4, we have

$$\begin{aligned}
I_U &= \int \int_U \int \frac{|g(s, X(s, t, x, v), V(s, t, x, v))|}{|X(s, t, x, v) - \hat{X}(s)|^2} dv dx ds \\
&= \int \int_U \int \frac{|F(V(s, t, x, v)) - f(s, X(s, t, x, v), V(s, t, x, v))|}{|X(s, t, x, v) - \hat{X}(s)|^2} dv dx ds \\
&= \int \int_U \int \frac{|-f(t, x, v)|}{|X(s, t, x, v) - \hat{X}(s)|^2} dv dx ds \\
&= \int_{t-\Delta}^t \int_{|v|>P \cap |v-\hat{V}(t)|>P} \int_{|X(s, t, x, v) - \hat{X}(s)|>R} \frac{f(t, x, v)}{|X(s, t, x, v) - \hat{X}(s)|^2} dx dv ds \\
&= \int_{|v|>P} \int f(t, x, v) \left(\int_{t-\Delta}^t |X(s, t, x, v) - \hat{X}(s)|^{-2} \chi_U(s, x, v) ds \right) dx dv \\
&\leq \frac{C}{RP} \int \int_{|v|>P} f(t, x, v) dv dx \\
&\leq \frac{C}{RP^3} \int \int_{|v|>P} |v|^2 f(t, x, v) dv dx
\end{aligned}$$

and so

$$I_U \leq \frac{C}{RP^3}. \quad (26)$$

Finally, collecting the estimates (22), (23), and (26), we find,

$$\frac{1}{\Delta} \int_{t-\Delta}^t |E(s, \hat{X}(s))| ds \leq C \left(P^{\frac{4}{3}} + RQ^3(t) + \frac{1}{\Delta RP^3} \right).$$

We take $R = Q^{-\frac{32}{15}}(t)$ and then

$$\frac{1}{\Delta} \int_{t-\Delta}^t |E(s, \hat{X}(s))| ds \leq CQ^{\frac{13}{15}}(t).$$

Using (III), we have

$$\begin{aligned}
\frac{1}{\Delta} \int_{t-\Delta}^t \left(|E(s, \hat{X}(s)) + A(s, \hat{X}(s))| \right) ds &\leq CQ^{\frac{13}{15}}(t) + \frac{1}{\Delta} \int_{t-\Delta}^t C^{(0)} R^{-2}(\hat{X}(s)) ds \\
&\leq CQ^{\frac{13}{15}}(t) + \frac{1}{\Delta} \int_{t-\Delta}^t C^{(0)} ds \\
&\leq CQ^{\frac{13}{15}}(t) + \left((C^{(0)})^{\frac{15}{13}} \right)^{\frac{13}{15}} \\
&\leq CQ^{\frac{13}{15}}(t) + Q^{\frac{13}{15}}(t) \\
&\leq CQ^{\frac{13}{15}}(t).
\end{aligned} \quad (27)$$

Finally, we use the argument in Section 4.5 of [4] to bound the velocity support, since the power of $Q(t)$ is less than one. This work will be explored in Appendix B. Thus, for all $t \in [0, T]$,

$$Q(t) \leq C, \quad (28)$$

and this implies bounds on $Q_f(t)$ and $Q_g(t)$ for all $t \in [0, T]$. Furthermore, if $V(s, t, x, v)$ satisfies $f_0(X(0, t, x, v), V(0, t, x, v)) \neq 0$, then

$$|V(s, t, x, v)| \leq C$$

for any $s \in [0, T]$, including $V(t, t, x, v) = v$.

Notice then, the bound on the velocity support implies a priori bounds on $\|\rho(t)\|_{L^\infty}$, $\|E(t)\|_{L^\infty}$, $\|\nabla_x E(t)\|_{L^\infty}$, and $\|\nabla_{x,v} f(t)\|_{L^\infty}$ for all $t \in [0, T]$.

Now that $\|\rho(t)\|_\infty \leq C$ and $\|\nabla \rho(t)\|_\infty \leq C$ for all $t \in [0, T]$, and $Q_f(T)$ is finite, we apply Lemma 4 since $p > 3$, and find $\|\rho(t)\|_p \leq C$ for all $t \in [0, T]$, and the proof of Theorem 1 is complete.

3 Section 3

To conclude the paper, this section contains the proofs of Lemmas 2 through 4.

Proof (Lemma 2) : Let $q, T > 0$ be given with $\|\rho(t)\|_\infty \leq C$ for all $t \in [0, T]$. Let $b \in [0, \frac{5}{18})$ be given with $b \leq \frac{2}{q}$. Consider $|x| \geq 1$ and define

$$\eta := (\|\rho(t)\|_q R^{-q}(x))^b$$

and divide the field into the following pieces :

$$|E(t, x)| \leq I + II + III.$$

where

$$\begin{aligned} I &:= \int_{|x-y| < \eta} |\rho(t, y)| |x-y|^{-2} dy, \\ II &:= \int_{\eta < |x-y| < \frac{1}{2}|x|} |\rho(t, y)| |x-y|^{-2} dy, \\ III &:= \int_{|x-y| > \frac{1}{2}|x|} |\rho(t, y)| |x-y|^{-2} dy. \end{aligned}$$

Then, the first estimate satisfies

$$I \leq C \|\rho(t)\|_\infty \eta. \quad (29)$$

The second estimate satisfies, for any $m \in [0, \frac{1}{3})$,

$$\begin{aligned} II &\leq C (\|\rho(t)\|_q R^{-q}(x))^m \int_{\eta < |x-y| < \frac{1}{2}|x|} |\rho(t, y)|^{1-m} |x-y|^{-2} dy \\ &\leq C (\|\rho(t)\|_q R^{-q}(x))^m \int_{\eta < |x-y| < \frac{1}{2}|x|} \left(k^{\frac{3(1-m)}{5}}(t, y) |x-y|^{-2} + k^{\frac{1-m}{2}}(t, y) |x-y|^{-2} \right) dy \\ &\leq C (\|\rho(t)\|_q R^{-q}(x))^m \left[\left(\int_{\eta < |x-y| < \frac{1}{2}|x|} |x-y|^{-\frac{4}{1+m}} dy \right)^{\frac{1+m}{2}} \right. \\ &\quad \left. + \left(\int_{\eta < |x-y| < \frac{1}{2}|x|} |x-y|^{-\frac{10}{3m+2}} dy \right)^{\frac{3m+2}{5}} \right] \\ &\leq C (\|\rho(t)\|_q R^{-q}(x))^m \left[\eta^{-2+3(\frac{1+m}{2})} + \eta^{-2+3(\frac{3m+2}{5})} \right] \\ &\leq C (\|\rho(t)\|_q R^{-q}(x))^m \begin{cases} \eta^{\frac{3m-1}{2}}, & \eta \geq 1 \\ \eta^{\frac{9m-4}{5}}, & \eta \leq 1 \end{cases} \end{aligned}$$

Then, for $\eta \geq 1$, we choose $m = \frac{3b}{2+3b}$, and for $\eta \leq 1$, we choose $m = \frac{9b}{5+9b}$. To guarantee convergence of the above integrals, we must have $m < \frac{1}{3}$, and thus, $b < \frac{5}{18}$. Thus, we find

$$II \leq C (\|\rho(t)\|_q R^{-q}(x))^b. \quad (30)$$

Finally, for $b \leq \frac{2}{q}$,

$$\begin{aligned} III &\leq \|\rho(t)\|_q^b \left(\frac{1}{2}|x|\right)^{-qb} \int_{|x-y| > \frac{1}{2}|x|} |\rho(t, y)|^{1-b} |x-y|^{bq-2} R^{-bq}(y) dy \\ &\leq C (\|\rho(t)\|_q R^{-q}(x))^b \int_{|x-y| > \frac{1}{2}|x|} |\rho(t, y)|^{1-b} \left(\frac{1}{4}R(y)\right)^{bq-2} R^{-bq}(y) dy \\ &\leq C (\|\rho(t)\|_q R^{-q}(x))^b \int_{|x-y| > \frac{1}{2}|x|} (k^{\frac{1-b}{2}}(t, y) + k^{\frac{3}{5}(1-b)}(t, y)) R^{-2}(y) dy \\ &\leq C (\|\rho(t)\|_q R^{-q}(x))^b \left[\left(\int R^{-\frac{4}{1+b}}(y) dy \right)^{\frac{1+b}{2}} + \left(\int R^{-\frac{10}{3b+2}}(y) dy \right)^{\frac{3b+2}{5}} \right] \\ &\leq C (\|\rho(t)\|_q R^{-q}(x))^b \end{aligned}$$

for $-\frac{4}{1+b} < -3$, which is satisfied since $b < \frac{5}{18}$.

Combining the estimates for I , II , and III , the lemma follows.

Proof (Lemma 3) : Let $q, T > 0$ be given with $\|\rho(t)\|_\infty \leq C$ and $\|\nabla\rho(t)\|_\infty \leq C$ for all $t \in [0, T]$. Let $a \in [0, 1)$ be given with $a \leq \frac{3}{q}$. Consider $|x| \geq 1$ and define

$$\eta := (\|\rho(t)\|_q R^{-q}(x))^a.$$

For any $i, k = 1, 2, 3$, we have

$$\begin{aligned} |\partial_{x_i} E_k(t, x)| &= |\int_{|x-y|<\eta} \partial_{y_i} \rho(t, y) \frac{(x-y)_k}{|x-y|^3} dy| + |\int_{|x-y|=\eta} \rho(t, y) \frac{(x-y)_k}{|x-y|^3} \frac{(x-y)_i}{|x-y|} dS_y| \\ &\quad + |\int_{|x-y|>\eta} \rho(t, y) \partial_{y_i} \left(\frac{(x-y)_k}{|x-y|^3} \right) dy| \\ &=: I + II + III. \end{aligned}$$

Then,

$$\begin{aligned} I &\leq \int_{|x-y|<\eta} \|\nabla_x \rho(t)\|_\infty |x-y|^{-2} dy \\ &\leq C\eta. \end{aligned}$$

We estimate II for the large $|x|$ and small $|x|$ cases. For $|x| > 2\eta$,

$$\begin{aligned} II &\leq \int_{|x-y|=\eta} (\|\rho(t)\|_q R^{-q}(y))^a |\rho(t, y)|^{1-a} |x-y|^{-2} dS_y \\ &\leq \|\rho(t)\|_q^a \|\rho(t)\|_{L^\infty}^{1-a} \eta^{-2} \int_{|x-y|=\eta} R^{-aq} (|x| - \eta) dS_y \\ &\leq C \|\rho(t)\|_q^a \eta^{-2} R^{-aq} \left(\frac{1}{2}|x| \right) \eta^2 \\ &\leq C (\|\rho(t)\|_q R^{-q}(x))^a. \end{aligned}$$

For $|x| < 2\eta$,

$$\begin{aligned} II &\leq \|\rho(t)\|_\infty \eta^{-2} \int_{|x-y|=\eta} dS_y \\ &\leq C \\ &\leq C(2\eta|x|^{-1}) \\ &\leq C\eta. \end{aligned}$$

Then,

$$\begin{aligned} III &\leq C \int_{\eta < |x-y| < \frac{1}{2}|x|} |\rho(t, y)| |x-y|^{-3} dy + C \int_{|x-y| > \frac{1}{2}|x|} |\rho(t, y)| |x-y|^{-3} dy \\ &=: A + B. \end{aligned}$$

Estimating A , we have for $n \in [0, 1)$,

$$\begin{aligned} A &\leq \|\rho(t)\|_q^n \int_{\eta < |x-y| < \frac{1}{2}|x|} |\rho(t, y)|^{1-n} R^{-nq}(y) |x-y|^{-3} dy \\ &\leq C \|\rho(t)\|_q^n R^{-nq}(\tfrac{1}{2}|x|) \int_{\eta < |x-y| < \frac{1}{2}|x|} (k^{\frac{1-n}{2}}(t, y) + k^{\frac{3(1-n)}{5}}(t, y)) |x-y|^{-3} dy \\ &\leq C(\|\rho(t)\|_q R^{-q}(x))^n \left[\left(\int_{\eta < |x-y| < \frac{1}{2}|x|} |x-y|^{-\frac{6}{1+n}} dy \right)^{\frac{1+n}{2}} \right. \\ &\quad \left. + \left(\int_{\eta < |x-y| < \frac{1}{2}|x|} |x-y|^{-\frac{15}{3n+2}} dy \right)^{\frac{3n+2}{5}} \right] \\ &\leq C(\|\rho(t)\|_q R^{-q}(x))^n \left[\left(\int_{\eta}^{\infty} r^{-\frac{6}{1+n}+2} dr \right)^{\frac{1+n}{2}} + \left(\int_{\eta}^{\infty} r^{-\frac{15}{3n+2}+2} dr \right)^{\frac{3n+2}{5}} \right] \\ &\leq C(\|\rho(t)\|_q R^{-q}(x))^n (\eta^{\frac{3}{2}(1+n)-3} + \eta^{\frac{3}{5}(3n+2)-3}) \\ &\leq C(\|\rho(t)\|_q R^{-q}(x))^n (\eta^{-\frac{3}{2}(1-n)} + \eta^{-\frac{9}{5}(1-n)}) \\ &\leq C(\|\rho(t)\|_q R^{-q}(x))^n \begin{cases} \eta^{-\frac{9}{5}(1-n)}, & \eta \leq 1 \\ \eta^{-\frac{3}{2}(1-n)}, & \eta \geq 1 \end{cases} \end{aligned}$$

For $\eta \leq 1$, we may choose $n = \frac{14a}{9a+5}$, and for $\eta \geq 1$, we may choose $n = \frac{5a}{3a+2}$. Thus,

$$A \leq C(\|\rho(t)\|_q R^{-q}(x))^a.$$

Finally, we estimate B and find

$$\begin{aligned} B &\leq \int_{|x-y| > \frac{1}{2}|x|} |\rho(t, y)| |x-y|^{-3} dy \\ &\leq \|\rho(t)\|_q^a (\tfrac{1}{2}|x|)^{-aq} \int_{|x-y| > \frac{1}{2}|x|} |\rho(t, y)|^{1-a} |x-y|^{aq-3} R^{-aq}(y) dy \\ &\leq C(\|\rho(t)\|_q R^{-q}(x))^a \int_{|x-y| > \frac{1}{2}|x|} (k^{\frac{1-a}{2}}(t, y) + k^{\frac{3}{5}(1-a)}(t, y)) R^{-3}(y) dy \\ &\leq C(\|\rho(t)\|_q R^{-q}(x))^a \left[\left(\int R^{-\frac{6}{1+a}}(y) dy \right)^{\frac{1+a}{2}} + \left(\int R^{-\frac{15}{3a+2}}(y) dy \right)^{\frac{3a+2}{5}} \right] \\ &\leq C(\|\rho(t)\|_q R^{-q}(x))^a \end{aligned}$$

for $-\frac{6}{1+a} < -3$, which is satisfied since $a < 1$.

Combining the estimates for I , II , A , and B , the lemma follows.

Proof (Lemma 4) : Let $T > 0$ and $q \in [0, \frac{54}{13})$ be given with $\|\rho(t)\|_\infty \leq C$ and $\|\nabla \rho(t)\|_\infty \leq C$ for all $t \in [0, T]$ and $Q_f(T) < \infty$. Let $t \in [0, T]$ be given. For any $D > 0$, taking $|x| \leq D$, we find

$$\begin{aligned} |\rho(t, x)| &\leq \|\rho(t)\|_\infty \\ &\leq \|\rho(t)\|_\infty \frac{R^q(D)}{R^q(x)} \\ &\leq CR^{-q}(x). \end{aligned} \tag{31}$$

Now, define $C^{(3)} := \max\{1, 8TQ_g(T)\}$ and let $|x| \geq C^{(3)}$. Define for every $t \in [0, T]$ and $x \in \mathbb{R}^3$,

$$\mathcal{E}(t, x) = E(t, x) + A(t, x).$$

From the Vlasov equation,

$$\frac{\partial}{\partial s}(g(s, X(s), V(s))) = -\mathcal{E}(s, X(s)) \cdot \nabla_v F(V(s)),$$

and thus

$$g(t, x, v) = g(0, X(0), V(0)) - \int_0^t \mathcal{E}(s, X(s)) \cdot \nabla_v F(V(s)) \, ds. \tag{32}$$

Thus, to estimate ρ , we must consider $\int \mathcal{E}(s, X(s)) \cdot \nabla_v F(V(s)) \, dv$. Assume f is nonzero along $(X(s), V(s))$. Then,

$$\begin{aligned} \left| \int \mathcal{E}(s, X(s)) \cdot \nabla_v F(V(s)) \, dv \right| &\leq \left| \int \mathcal{E}(s, X(s)) \cdot (\nabla_v F(V(s)) - \nabla_v F(v + \int_s^t \mathcal{E}(\tau, x) \, d\tau)) \, dv \right| \\ &\quad + \left| \int (\mathcal{E}(s, X(s)) - \mathcal{E}(s, x + (s-t)v)) \right. \\ &\quad \left. \cdot \nabla_v F(v + \int_s^t \mathcal{E}(\tau, x) \, d\tau) \, dv \right| \\ &\quad + \left| \int \nabla_v \cdot (F(v) \mathcal{E}(s, x + (s-t)v)) \, dv \right| \\ &\quad + \left| \int F(v) \nabla_v \cdot (\mathcal{E}(s, x + (s-t)v)) \, dv \right| \\ &=: I + II + III + IV. \end{aligned}$$

Using Lemmas 2 and 3, as well as (III), we find $a \in [0, 1)$ and $b \in [0, \frac{5}{18})$ with $a \leq \frac{3}{q}$, $b \leq \frac{2}{q}$, and $a + b = 1$ such that

$$|\mathcal{E}(t, x)| \leq C (\|\rho(t)\|_q R^{-q}(x))^b$$

and

$$|\nabla \mathcal{E}(t, x)| \leq C (\|\rho(t)\|_q R^{-q}(x))^a.$$

By the Mean Value Theorem, for $\tau \in [s, t]$ and $i = 1, 2, 3$, there exist ξ_1^i on the line segment between $X(\tau)$ and x such that

$$\mathcal{E}_i(\tau, X(\tau)) - \mathcal{E}_i(\tau, x) = \nabla_x \mathcal{E}_i(\tau, \xi_1^i) \cdot (X(\tau) - x).$$

Hence,

$$\begin{aligned} I &\leq \int_{|v| \leq Q_g(T)} |\mathcal{E}(s, X(s))| \|\nabla^2 F\|_\infty |V(s) - \left(v + \int_s^t \mathcal{E}(\tau, x) d\tau\right)| dv \\ &\leq C \int_{|v| \leq Q_g(T)} (\|\rho(s)\|_q R^{-q}(X(s)))^b \left| \int_s^t (\mathcal{E}(\tau, X(\tau)) - \mathcal{E}(\tau, x)) d\tau \right| dv \\ &\leq C \int_{|v| \leq Q_g(T)} (\|\rho(s)\|_q R^{-q}(X(s)))^b \int_s^t \sup_i |\nabla \mathcal{E}(\tau, \xi_1^i)| |X(\tau) - x| d\tau dv \\ &\leq C \int_{|v| \leq Q_g(T)} (\|\rho(s)\|_q R^{-q}(X(s)))^b T Q_g(T) \int_s^t \sup_i (\|\rho(\tau)\|_q R^{-q}(\xi_1^i))^a d\tau dv. \end{aligned}$$

Since we know, by (11), for any $i = 1, 2, 3$

$$\begin{aligned} |\xi_1^i| &\geq |X(\tau)| - |X(\tau) - x| \\ &\geq \frac{1}{2}|x| - T Q_g(T) \\ &\geq \frac{1}{4}|x|, \end{aligned}$$

we find

$$\begin{aligned} I &\leq C \left(\|\rho(s)\|_q R^{-q}(\tfrac{1}{2}|x|) \right)^b R^{-aq}(\tfrac{1}{4}|x|) \int_s^t \|\rho(\tau)\|_q^a d\tau \\ &\leq C R^{-q(a+b)}(x) \left(\|\rho(s)\|_q^b \int_s^t \|\rho(\tau)\|_q^a d\tau \right). \end{aligned}$$

Similarly, using the above lemmas and the Mean Value Theorem, for any $i = 1, 2, 3$ there exist ξ_2^i between $X(\tau)$ and $x + (s - t)v$ such that

$$\begin{aligned} II &\leq C \sum_{i=1}^3 \int_{|v| \leq Q_g(T)} \sup_i |\nabla \mathcal{E}(s, \xi_2^i)| |X(s) - (x + (s - t)v)| \|\nabla F\|_\infty dv \\ &\leq C \sum_{i=1}^3 \int_{|v| \leq Q_g(T)} \sup_i (\|\rho(s)\|_q R^{-q}(\xi_2^i))^a \left| \int_s^t \int_\tau^t \mathcal{E}(\iota, X(\iota)) d\iota d\tau \right| dv. \end{aligned}$$

Since we know, using (11), for any $i = 1, 2, 3$

$$\begin{aligned} |\xi_2^i| &\geq |X(s)| - |X(s) - (x + (s - t)v)| \\ &\geq \frac{1}{2}|x| - \left| \int_s^t (V(\tau) - v) d\tau \right| \\ &\geq \frac{1}{2}|x| - 2TQ_g(T) \\ &\geq \frac{1}{4}|x|, \end{aligned}$$

it follows that

$$\begin{aligned} II &\leq C \int_{|v| \leq Q_g(T)} \left(\|\rho(s)\|_q R^{-q}(\tfrac{1}{4}|x|) \right)^a \int_s^t (\|\rho(s)\|_q R^{-q}(X(\tau)))^b d\tau dv \\ &\leq CR^{-q(a+b)}(x) \left(\|\rho(s)\|_q^a \int_s^t \|\rho(\tau)\|_q^b d\tau \right). \end{aligned}$$

By the Divergence Theorem,

$$III = 0,$$

and finally,

$$\begin{aligned} IV &\leq \int_{|v| \leq Q_g(T)} \|F\|_\infty |\nabla_x \cdot \mathcal{E}(s, x + (s - t)v)| |s - t| dv \\ &\leq C \int_{|v| \leq Q_g(T)} |\rho(s, x + (s - t)v)| dv \end{aligned}$$

Collecting the estimates for $I - IV$, we have

$$\begin{aligned}
|\int \mathcal{E}(s, X(s)) \cdot \nabla_v F(V(s)) dv| &\leq C \left(R^{-q(a+b)}(x) (\|\rho(s)\|_q^b \int_s^t \|\rho(\tau)\|_q^a d\tau) \right. \\
&\quad \left. + R^{-q(a+b)}(x) (\|\rho(s)\|_q^a \int_s^t \|\rho(\tau)\|_q^b d\tau) \right. \\
&\quad \left. + \int_{|v| \leq Q_g(T)} |\rho(s, x + (s-t)v)| dv \right). \tag{33}
\end{aligned}$$

Since $a + b = 1$, we proceed from (32) and using (33), (IV), and (28), we find

$$\begin{aligned}
|\rho(t, x)| &= \left| \int g(t, x, v) dv \right| \\
&\leq \int_{|v| \leq Q_g(T)} |g(0, X(0), V(0))| dv + \left| \int_0^t \int_{|v| \leq Q_g(T)} \mathcal{E}(s, X(s)) \cdot \nabla_v F(V(s)) dv ds \right| \\
&\leq CR^{-q}(x) + C \int_0^t \left(R^{-q(a+b)}(x) (\|\rho(s)\|_q^b \int_s^t \|\rho(\tau)\|_q^a d\tau) \right. \\
&\quad \left. + R^{-q(a+b)}(x) (\|\rho(s)\|_q^a \int_s^t \|\rho(\tau)\|_q^b d\tau) + \int_{|v| \leq Q_g(T)} |\rho(s, x + (s-t)v)| dv \right) ds \\
&\leq CR^{-q}(x) \left(1 + \int_0^t \|\rho(s)\|_q^b \int_s^t \|\rho(\tau)\|_q^a d\tau ds + \int_0^t \|\rho(s)\|_q^a \int_s^t \|\rho(\tau)\|_q^b d\tau ds \right) \\
&\quad + C \int_{|v| \leq Q_g(T)} \int_0^t |\rho(s, x + (s-t)v)| ds dv \\
&\leq CR^{-q}(x) \left(1 + \left(\int_0^t \|\rho(s)\|_q^b ds \right) \left(\int_0^t \|\rho(\tau)\|_q^a d\tau \right) + \left(\int_0^t \|\rho(s)\|_q^a ds \right) \left(\int_0^t \|\rho(\tau)\|_q^b d\tau \right) \right) \\
&\quad + C \int_{|v| \leq Q_g(T)} \int_0^t |\rho(s, x + (s-t)v)| ds dv.
\end{aligned}$$

Then, applying Hölder's inequality twice,

$$\begin{aligned}
|\rho(t, x)| &\leq CR^{-q}(x) \left(1 + 2T^2 \left(\int_0^t \|\rho(s)\|_q ds \right)^{a+b} \right. \\
&\quad \left. + R^q(x) \int_{|v| \leq Q_g(T)} \int_0^t \|\rho(s)\|_q R^{-q}(x + (s-t)v) ds dv \right) \\
&\leq CR^{-q}(x) \left(1 + \int_0^t \|\rho(s)\|_q ds + R^q(x) R^{-q}(\frac{7}{8}|x|) Q_g^3(T) \int_0^t \|\rho(s)\|_q ds \right) \\
&\leq CR^{-q}(x) \left(1 + \int_0^t \|\rho(s)\|_q ds \right).
\end{aligned}$$

Combining this with (31) (with $D = C^{(3)}$) and multiplying by $R^q(x)$, we have for all x ,

$$R^q(x)|\rho(t, x)| \leq C(1 + \int_0^t \|\rho(s)\|_q ds).$$

Since the right side of the inequality is independent of x , we take the supremum over all x to find,

$$\|\rho(t)\|_q \leq C(1 + \int_0^t \|\rho(s)\|_q ds)$$

and by Gronwall's Inequality,

$$\|\rho(t)\|_q \leq C.$$

Thus, $\|\rho(t)\|_q$ is bounded for all $t \in [0, T]$. Notice, too, that the choice of $a + b = 1$ forces $q < \frac{54}{13}$, and the proof of the lemma is complete.

A Appendix A

In this appendix, we will explore the argument used in Section 4.2.6 of [4] to bound derivatives of f and E .

Let D be any x derivative. Then, using the Vlasov equation,

$$\partial_t(Df) + v \cdot \nabla_x(Df) - E \cdot \nabla_v(Df) = DE \cdot \nabla_v f$$

so that

$$\frac{d}{ds} Df(s, X(s), V(s)) = DE \cdot \nabla_v f(s, X(s), V(s)).$$

Hence,

$$|Df(t, x, v)| \leq |Df(0, X(0), V(0))| + \int_0^t |DE \cdot \nabla_v f(s, X(s), V(s))| ds.$$

Define

$$|f(s)|_1 = \sup_{x,v} |\partial_x f(s, x, v)| + \sup_{x,v} |\partial_v f(s, x, v)|$$

and

$$|E(s)|_1 = \sup_x |\partial_x E(s, x)|.$$

Then,

$$|Df(t, x, v)| \leq C + \int_0^t |E(s)|_1 |f(s)|_1 ds.$$

We see that $\partial_v f$ satisfies an inequality of similar type because

$$\partial_t(\partial_v f) + v \cdot \nabla_x(\partial_v f) - E \cdot \nabla_v(\partial_v f) = -\partial_v v \cdot \nabla_x f.$$

It follows that

$$|f(t)|_1 \leq C + \int_0^t (1 + |E(s)|_1) |f(s)|_1 ds.$$

However, from (18), we can conclude (with $d = \|\nabla_x \rho\|_\infty$) that

$$|E(s)|_1 \leq C(1 + \ln^* |f(s)|_1)$$

where

$$\ln^* s = \begin{cases} s & 0 \leq s \leq 1 \\ 1 + \ln s & s > 1. \end{cases}$$

Therefore, for $t \leq T$,

$$|f(t)|_1 \leq C \left(1 + \int_0^t (1 + \ln^* |f(s)|_1) |f(s)|_1 ds \right).$$

It now follows by an application of Gronwall's Inequality that

$$|f(t)|_1 \leq C$$

and

$$|E(t)|_1 \leq C$$

for $t \leq T$. This concludes the argument to bound field derivatives and derivatives of the density, and thus ends Appendix A.

B Appendix B

The bound on $Q(t)$ is obtained as follows. From (8) and (27), we have

$$\begin{aligned} |\hat{V}(t)| &\leq |\hat{V}(t - \Delta)| + \int_{t-\Delta}^t |E(s, \hat{X}(s))| ds \\ &\leq Q(t - \Delta) + C\Delta Q^{\frac{13}{15}}(t). \end{aligned}$$

Since the above constant is independent of the particular characteristic, we find

$$Q(t) \leq Q(t - \Delta) + C\Delta Q^{\frac{13}{15}}(t)$$

for

$$\begin{aligned}\Delta &= \min \left\{ t, \frac{1}{4C^{(2)}} Q^{-\frac{4}{3}}(t) \cdot Q^{\frac{13}{20}}(t) \right\}. \\ &= \min \left\{ t, \frac{1}{4C^{(2)}} Q^{-\frac{41}{60}}(t) \right\}\end{aligned}$$

Since Q is non-decreasing, there exists T_1 such that

$$\Delta = \begin{cases} t & t \leq T_1 \\ \frac{1}{4C^{(2)}} Q^{-\frac{41}{60}}(t) & t \geq T_1. \end{cases}$$

Take t_0 in the interval of existence. Without loss of generality, $t_0 > T_1$. Let

$$t_1 = t_0 - \frac{1}{4C^{(2)}} Q^{-\frac{41}{60}}(t_0),$$

$$t_{i+1} = t_i - \frac{1}{4C^{(2)}} Q^{-\frac{41}{60}}(t_i) \quad (i = 1, 2, \dots)$$

as long as $t_i > T_1$. Then

$$t_i - t_{i+1} = \frac{1}{4C^{(2)}} Q^{-\frac{41}{60}}(t_i) \geq \frac{1}{4C^{(2)}} Q^{-\frac{41}{60}}(t_0)$$

which is a uniform lower bound on the length of each subinterval. So, there is a first i , say $i = k$, such that $t_k \leq T_1$. Thus, $t_k \geq 0$ and therefore

$$(t_0 - t_1) + (t_1 - t_2) + \dots + (t_{k-1} - t_k) \geq k \cdot \frac{1}{4C^{(2)}} Q^{-\frac{41}{60}}(t_0)$$

which implies that

$$Q^{-\frac{41}{60}}(t_0) \cdot k \leq 4C^{(2)} t_0.$$

Now we have

$$\begin{aligned}Q(t_0) &= Q(t_k) + \sum_{i=0}^{k-1} [Q(t_i) - Q(t_{i+1})] \\ &\leq Q(t_k) + C \sum_{i=0}^{k-1} \Delta \cdot Q^{\frac{13}{15}}(t_i) \\ &\leq Q(T_1) + C \sum_{i=0}^{k-1} \frac{1}{4C^{(2)}} Q^{-\frac{41}{60}}(t_0) \cdot Q^{\frac{13}{15}}(t_i) \\ &\leq Q(T_1) + C \cdot k Q^{-\frac{41}{60}}(t_0) \cdot Q^{\frac{13}{15}}(t_0) \\ &\leq C t_0 Q^{\frac{13}{15}}(t_0).\end{aligned}$$

Therefore, $Q(t_0)$ is bounded, and the proof is complete. This ends Appendix B.

References

- [1] Batt, J. Global symmetric solutions of the initial-value problem of stellar dynamics. *J. Diff. Eq.* **1977**, 25 : 342-364.
- [2] Batt, J.; Rein, G. Global classical solutions of the periodic Vlasov-Poisson system in three dimensions. *C. R. Academy of Sci.* **1991**, 313(1): 411-416.
- [3] Caglioti, E.; Caprino, S.; Marchioro, C.; Pulvirenti, M. The Vlasov equation with infinite charge. *Arch. Ration. Mech. Anal.* **2001**, 159:85-108.
- [4] Glassey, R. *The Cauchy Problem in Kinetic Theory*; S.I.A.M: Philadelphia, 1996.
- [5] Glassey, R.; Strauss, W. Singularity formation in a collisionless plasma could occur only at high velocities. *Arch. Ration. Mech. Ana.* **1986**, 92:59-90.
- [6] Horst, E. On the classical solutions of the initial value problem for the unmodified nonlinear Vlasov-equation, part I. *Math. Methods Appl. Sci.* **1981**, 3:229-248.
- [7] Horst, E. On the classical solutions of the initial value problem for the unmodified nonlinear Vlasov-equation, part II. *Math. Methods Appl. Sci.* **1982**, 4:19-32.
- [8] Horst, E. On the asymptotic growth of the solutions of the Vlasov-Poisson system. *Math. Methods Appl. Sci.* **1993**, 16:75-85.
- [9] Jabin, P. E. The Vlasov-Poisson system with infinite charge and energy. *J. Statist. Phys.* **2001**, 103(5/6):1107-1123.
- [10] Kurth, R. Das anfangswertproblem der stellardynamik. *Z. Astrophys.* **1952** 30:213-229.
- [11] Lions, P.L.; Pertham, B. Propagation of moments and regularity for the three dimensional Vlasov-Poisson system. *Invent. Math.* **1991**, 105:415-430.
- [12] Okabe, S.; Ukai, T. On classical solutions in the large in time of two-dimensional Vlasov's equation. *Osaka J. Math.* **1978**, 15:245-261.
- [13] Pankavich, S. Global existence for the Radial Vlasov-Poisson System with Steady Spatial Asymptotics. **2004** (submitted for publication)
- [14] Pfaffelmoser, K. Global classical solution of the Vlasov-Poisson system in three dimensions for general initial data. *J. Diff. Eq.* **1992**, 95(2):281-303.
- [15] Rein, G.; Rendall, A. Global existence of classical solutions to the Vlasov-Poisson system in a three dimensional cosmological setting. *Arch. Ration. Mech. Anal.* **1994** 126:183-201.

- [16] Schaeffer, J. Global existence of smooth solutions to the Vlasov-Poisson system in three dimensions. *Comm. PDE.* **1991**, 16(8/9):1313-1335.
- [17] Schaeffer, J. Steady spatial asymptotics for the Vlasov-Poisson system. *Math. Methods Appl. Sci.* **2003**, 26:273-296.
- [18] Schaeffer, J. The Vlasov-Poisson system with steady spatial asymptotics. *Comm. PDE.* **2003**, 28(5/6):1057-1084.
- [19] Wollman, S. Global-in-time solutions of the two dimensional Vlasov-Poisson system. *Comm. Pure Appl. Math.* **1980**, 33:173-197.